

Recent results from nuclear quantum Monte Carlo

Robert B. Wiringa, Physics Division, Argonne National Laboratory

Ivan Brida, Los Alamos
Joseph Carlson, Los Alamos
Kenneth M. Nollett, Ohio
Saori Pastore, Argonne
Steven C. Pieper, Argonne
Rocco Schiavilla, JLab & ODU

WORK NOT POSSIBLE WITHOUT EXTENSIVE COMPUTER RESOURCES

Argonne Laboratory Computing Resource Center (Jazz & Fusion)
Argonne Math. & Comp. Science Division (BlueGene/L & SiCortex)
Argonne Leadership Computing Facility (BlueGene/P)



Physics Division

Work supported by U.S. Department
of Energy, Office of Nuclear Physics

Ab Initio CALCULATIONS OF LIGHT NUCLEI

GOALS

Understand nuclei at the level of elementary interactions between individual nucleons, including

- Binding energies, excitation spectra, relative stability
- Densities, electromagnetic moments, transition amplitudes, cluster-cluster overlaps
- Low-energy NA & AA' scattering, asymptotic normalizations, astrophysical reactions

REQUIREMENTS

- Two-nucleon potentials that accurately describe elastic NN scattering data
- Consistent three-nucleon potentials and electroweak current operators
- Accurate methods for solving the many-nucleon Schrödinger equation

RESULTS

- Quantum Monte Carlo methods can evaluate realistic Hamiltonians accurate to $\sim 1-2\%$
- About 100 states calculated for $A \leq 12$ nuclei in good agreement with experiment
- Applications to elastic & inelastic e, π scattering, $(e, e'p)$, (d, p) reactions, etc.
- Electromagnetic moments, $M1$, $E2$, F, GT transitions calculated
- ${}^5\text{He} = n\alpha$ scattering and $3 \leq A \leq 9$ ANCs and widths

NUCLEAR HAMILTONIAN

$$H = \sum_i K_i + \sum_{i<j} v_{ij} + \sum_{i<j<k} V_{ijk}$$

$$K_i = -\frac{\hbar^2}{4} \left[\left(\frac{1}{m_p} + \frac{1}{m_n} \right) + \left(\frac{1}{m_p} - \frac{1}{m_n} \right) \tau_{iz} \right] \nabla_i^2$$

Argonne v18

$$v_{ij} = v_{ij}^\gamma + v_{ij}^\pi + v_{ij}^I + v_{ij}^S = \sum v_p(r_{ij}) O_{ij}^p$$

v_{ij}^γ : pp, pn & nn electromagnetic terms

$$v_{ij}^\pi \sim [Y_\pi(r_{ij}) \sigma_i \cdot \sigma_j + T_\pi(r_{ij}) S_{ij}] \otimes \tau_i \cdot \tau_j$$

$$v_{ij}^I = \sum_p I^p T_\pi^2(r_{ij}) O_{ij}^p$$





$$v_{ij}^S = \sum_p [P^p + Q^p r + R^p r^2] W(r) O_{ij}^p$$

$$\begin{aligned} O_{ij}^p &= [1, \sigma_i \cdot \sigma_j, S_{ij}, \mathbf{L} \cdot \mathbf{S}, \mathbf{L}^2, \mathbf{L}^2(\sigma_i \cdot \sigma_j), (\mathbf{L} \cdot \mathbf{S})^2] \\ &+ [1, \sigma_i \cdot \sigma_j, S_{ij}, \mathbf{L} \cdot \mathbf{S}, \mathbf{L}^2, \mathbf{L}^2(\sigma_i \cdot \sigma_j), (\mathbf{L} \cdot \mathbf{S})^2] \otimes \tau_i \cdot \tau_j \\ &+ [1, \sigma_i \cdot \sigma_j, S_{ij}, \mathbf{L} \cdot \mathbf{S}] \otimes T_{ij} \\ &+ [1, \sigma_i \cdot \sigma_j, S_{ij}, \mathbf{L} \cdot \mathbf{S}] \otimes (\tau_i + \tau_j)_z \end{aligned}$$

$$S_{ij} = 3\sigma_i \cdot \hat{r}_{ij} \sigma_j \cdot \hat{r}_{ij} - \sigma_i \cdot \sigma_j \quad T_{ij} = 3\tau_{iz} \tau_{jz} - \tau_i \cdot \tau_j$$


Wiringa, Stoks, & Schiavilla, PRC **51**, 38 (1995)





Argonne v_{18} fits Nijmegen PWA93 data base of 1787 pp & 2514 np observables for $E_{lab} \leq 350$ MeV with $\chi^2/\text{datum} = 1.1$ plus nn scattering length & ^2H binding energy

Argonne v_{18}



Uses 42 I^P , P^P , Q^P , R^P parameters [constrained so that $v_t(r=0) = 0$ & $\frac{\partial v_{p \neq t}}{\partial r} \Big|_{r=0} = 0$]
 plus $f_{\pi NN}$ coupling strength & one cutoff parameter in $Y_\pi(r)$, $T_\pi(r)$.


THREE-NUCLEON POTENTIALS

Urbana $V_{ijk} = V_{ijk}^{2\pi P} + V_{ijk}^R$



Carlson, Pandharipande, & Wiringa, NP **A401**, 59 (1983)

Illinois $V_{ijk} = V_{ijk}^{2\pi P} + V_{ijk}^{2\pi S} + V_{ijk}^{3\pi\Delta R} + V_{ijk}^R$



Pieper, Pandharipande, Wiringa, & Carlson, PRC **64**, 014001 (2001)

Illinois-7 has 4 strength parameters fit to 23 energy levels in $A \leq 10$ nuclei.

In light nuclei we find (thanks to large cancellation between $\langle K \rangle$ & $\langle v_{ij} \rangle$):

$$\langle V_{ijk} \rangle \sim (0.02 \text{ to } 0.07) \langle v_{ij} \rangle \sim (0.15 \text{ to } 0.5) \langle H \rangle$$

We expect $\langle \mathcal{V}_{ijkl} \rangle \sim 0.05 \langle V_{ijk} \rangle \sim (0.01 \text{ to } 0.03) \langle H \rangle \sim 1 \text{ MeV in } ^{12}\text{C} .$

VARIATIONAL MONTE CARLO

Minimize expectation value of H

$$E_V = \frac{\langle \Psi_V | H | \Psi_V \rangle}{\langle \Psi_V | \Psi_V \rangle} \geq E_0$$

using Metropolis Monte Carlo and trial function

$$|\Psi_V\rangle = \left[\mathcal{S} \prod_{i<j} (1 + U_{ij} + \sum_{k \neq i,j} U_{ijk}) \right] \left[\prod_{i<j} f_c(r_{ij}) \right] |\Phi_A(JMTT_3)\rangle$$

- single-particle $\Phi_A(JMTT_3)$ is fully antisymmetric and translationally invariant
- central pair correlations $f_c(r)$ keep nucleons at favorable pair separation
- pair correlation operators $U_{ij} = \sum_p u_p(r_{ij}) O_{ij}^p$ reflect influence of v_{ij}
- triple correlation operator U_{ijk} added when V_{ijk} is present
- multiple J^π states constructed and diagonalized for p-shell nuclei
- ability to construct clusterized or asymptotically correct trial functions

Ψ_V are spin-isospin vectors in $3A$ dimensions with $\sim 2^A \binom{A}{Z}$ components

Lomnitz-Adler, Pandharipande, & Smith, NP **A361**, 399 (1981)

Wiringa, PRC **43**, 1585 (1991)

GREEN'S FUNCTION MONTE CARLO

Projects out lowest energy state from variational trial function

$$\Psi(\tau) = \exp[-(H - E_0)\tau]\Psi_V = \sum_n \exp[-(E_n - E_0)\tau]a_n\psi_n$$
$$\Psi(\tau \rightarrow \infty) = a_0\psi_0$$

Evaluation of $\Psi(\tau)$ done stochastically in small time steps $\Delta\tau$

$$\Psi(\mathbf{R}_n, \tau) = \int G(\mathbf{R}_n, \mathbf{R}_{n-1}) \cdots G(\mathbf{R}_1, \mathbf{R}_0) \Psi_V(\mathbf{R}_0) d\mathbf{R}_{n-1} \cdots d\mathbf{R}_0$$

Mixed estimates used for expectation values

$$\langle O(\tau) \rangle = \frac{\langle \Psi(\tau) | O | \Psi(\tau) \rangle}{\langle \Psi(\tau) | \Psi(\tau) \rangle} \approx \langle O(\tau) \rangle_{\text{Mixed}} + [\langle O(\tau) \rangle_{\text{Mixed}} - \langle O \rangle_V]$$
$$\langle O(\tau) \rangle_{\text{Mixed}} = \frac{\langle \Psi_V | O | \Psi(\tau) \rangle}{\langle \Psi_V | \Psi(\tau) \rangle} \quad ; \quad \langle H(\tau) \rangle_{\text{Mixed}} = \frac{\langle \Psi(\tau/2) | H | \Psi(\tau/2) \rangle}{\langle \Psi(\tau/2) | \Psi(\tau/2) \rangle} \geq E_0$$


- Cannot propagate p^2 , L^2 , or $(\mathbf{L} \cdot \mathbf{S})^2$ operators \Rightarrow use $H' = AV8' + \tilde{V}_{ijk}$
- Fermion sign problem would limit maximum τ , but ...
- **Constrained-path propagation** removes steps that have $\overline{\Psi^\dagger(\tau, \mathbf{R})\Psi_V(\mathbf{R})} = 0$
- Multiple excited states of same J^π stay orthogonal

Pudliner, Pandharipande, Carlson, Pieper, & Wiringa, PRC **56**, 1720 (1997)


Wiringa, Pieper, Carlson, & Pandharipande, PRC **62**, 014001 (2000)

Pieper, Wiringa, & Carlson, PRC **70**, 054325 (2004)


EXAMPLES OF GFMC PROPAGATION



- Curve has $\sum_i a_i \exp(-E_i \tau)$ with $E_i = 1480, 340 \text{ \& } 20.2 \text{ MeV}$ (20.2 MeV is first ${}^4\text{He } 0^+$ excitation)
- Ψ_V has small amounts of 1.5 GeV contamination



- g.s. (1^+) & 3^+ stable after $\tau = 0.2 \text{ MeV}^{-1}$
- 2^+ (a broad resonance) never stable – decaying to separated α & d
- $E(\tau=0.2)$ is best GFMC estimate of resonance energy



Argonne v_{18} with Illinois-7 GFMC Calculations


Excitation energy (MeV)

Including IL7 gives
 • correct s.-o. splitting & ^{10}B g.s.

Legend for GFMC calculations:

- AV18 (Blue)
- AV18 + IL7 (Red)
- Expt. (Green)

States shown in legend: $7/2^+$, $5/2^+$, 0^+ , 4^+ , 3^+ , 1^+ , 2^+



1st AND 2nd (HOYLE) 0⁺ STATES IN ¹²C

Constructing the Jastrow part of the trial wave function is major effort:

- There are 5 *LS*-basis $J=0^+$ states in ¹²C in the 0*P* shell:
 $^1S[444], ^3P[4431], ^1S[4422], ^5D[4422], ^3P[4332]$
- All can be constructed by projections from a closed $(p3/2)^8$ shell (Carlson)
- Dominant $3-\alpha$ symmetry is easily constructed with one α in the 0*S* shell and two α s in the 0*P* shell (Pandharipande)
- Additional components generated by promoting one whole α to the 1*S*-0*D* shell, and also promoting pairs, e.g., 0*P*²0*D*² and 0*P*²1*S*²
- Total of 11 Jastrow components (some with considerable overlap) to be diagonalized

Challenge in GFMC propagation is keeping the 2nd state orthogonal to the ground state


UNEDF SciDAC grant to develop
general-purpose load-balancing library
(ADLB) to run under MPI on 32,768 nodes
with OpenMP for 4 cores/node

INCITE grant of Argonne's IBM
BlueGene/P time used for calculations





1st AND 2nd (HOYLE) 0⁺ STATES IN ¹²C – PRELIMINARY

Convergence as a function of imaginary time (τ)



	g.s. energy			2 nd 0 ⁺ E^*		
	VMC	GFMC	Expt.	VMC	GFMC	Expt.
AV18	-44.9(2)	-73.2(5)		10.0(3)	7.9(6)	
AV18+IL7	-65.7(2)	-93.3(4)	-92.16	14.7(2)	10.4(5)	7.65

1st AND 2nd (HOYLE) 0⁺ STATES IN ¹²C – PRELIMINARY





Central density dip for ground state may be interpreted as three α's in a triangle

Central density peak for Hoyle state may be evidence for a linear configuration of three α's

SINGLE-NUCLEON DENSITIES

$$\rho_{p,n}(r) = \sum_i \langle \Psi | \delta(r - r_i) \frac{1 \pm \tau_i}{2} | \Psi \rangle$$



RMS radii

	r_n	r_p	r_c	Expt
${}^4\text{He}$	1.45(1)	1.45(1)	1.67(1)	1.681(4)*
${}^6\text{He}$	2.86(6)	1.92(4)	2.06(4)	2.060(8)†
${}^8\text{He}$	2.79(3)	1.82(2)	1.94(2)	1.959(16)‡

*Sick, PRC **77**, 041302(R) (2008)


†Wang, *et al.*, PRL **93**, 142501 (2004)

‡Mueller, *et al.*, PRL **99**, 252501 (2007)

Brodeur, *et al.*, PRL **108**, 052504 (2012)

TWO-NUCLEON DENSITIES

$$\rho_{pp}(r) = \sum_{i < j} \langle \Psi | \delta(r - |\mathbf{r}_i - \mathbf{r}_j|) \frac{1 + \tau_i}{2} \frac{1 + \tau_j}{2} | \Psi \rangle$$



RMS radii

	r_{pp}	r_{np}	r_{nn}
${}^4\text{He}$	2.41	2.35	2.41
${}^6\text{He}$	2.51	3.69	4.40
${}^8\text{He}$	2.52	3.58	4.37

M1, E2, F, GT transitions

NO EFFECTIVE CHARGES!



$$E2 = e \sum_k \frac{1}{2} [r_k^2 Y_2(\hat{r}_k)] (1 + \tau_{kz})$$

$$M1 = \mu_N \sum_k [(L_k + g_p S_k)(1 + \tau_{kz})/2 + g_n S_k (1 - \tau_{kz})/2]$$

$$F = \sum_k \tau_{k\pm} ; \text{ GT} = \sum_k \sigma_k \tau_{k\pm}$$

Pervin, Pieper & Wiringa, PRC **76**, 064319 (2007)


Marcucci, Pervin, *et al.*, PRC **78**, 065501 (2008)



MAGNETIC MOMENTS w/ χ EFT EXCHANGE CURRENTS

Hybrid calculations using AV18+IL7 wave functions and χ EFT exchange currents developed in:


Pastore, Schiavilla, & Goity, PRC **78**, 064002 (2008) ; Pastore, *et al.*, PRC **80**, 034004 (2009)



M1 TRANSITIONS w/ χ EFT

- dominant contribution is from OPE
- five LECs at N3LO
- d_2^V and d_1^V are fixed assuming Δ resonance saturation
- d^S and c^S are fit to experimental μ_d and $\mu_S(^3\text{H}/^3\text{He})$
- c^V is fit to experimental $\mu_V(^3\text{H}/^3\text{He})$
- $\Lambda = 600$ MeV



Pastore, Pieper, Schiavilla & Wiringa (in preparation)



PRECISE EXPERIMENTAL TESTS OF THE ELECTROMAGNETIC RATES

New measurements of the lifetimes of the two ^{10}Be 2^+ states were made using the Doppler shift attenuation method following the $^7\text{Li}(^7\text{Li},\alpha)^{10}\text{Be}$ reaction. The $B(E2 \downarrow) = 9.2(3)e^2\text{fm}^4$ for the $J^\pi = 2_1^+$ state and $0.11(2)e^2\text{fm}^4$ for the $J^\pi = 2_2^+$ state.


A subsequent measurement of the lifetime of the 2_1^+ state in ^{10}C following the $p(^{10}\text{B},n)^{10}\text{C}$ reaction, got $B(E2 \downarrow) = 8.8(3)e^2\text{fm}^4$.



GFMC calculations using AV18 without or with IL2 or IL7 all get the 2_1^+ transition in ^{10}Be about right, but give widely varying predictions in ^{10}C . The latter appears much more sensitive to precise mixing of different symmetry state contributions, and thus to details of V_{ijk} .

TWO-NUCLEON HALO DENSITIES

$$\rho_{nn}(r) = \sum_{i < j} \langle \Psi(J^\pi, T, T_z = +1) | \delta(r - |\mathbf{r}_i - \mathbf{r}_j|) \tau_i^+ \tau_j^+ | \Psi(J^\pi, T, T_z = -1) \rangle$$



APPLICATIONS TO LIGHT-ION REACTIONS

The availability of radioactive-ion beams has renewed interest in reactions like (d,p) in inverse kinematics

We have helped analyze a number of RIB experiments such as $d(^8\text{Li},p)^9\text{Li}$ (ATLAS) & $d(^9\text{Li},t)^8\text{Li}$ (TRIUMF)


- PTOLEMY DWBA calculations for transfer
- (d,p) vertex from AV18
- (d,t) , $(^8\text{Li},^9\text{Li})$, etc. vertices computed as A -body overlaps using VMC
 $\langle \Psi_V(A-1) | a | \Psi_V(A) \rangle$
- Norm is spectroscopic factor
- Absolute prediction for $d\sigma/d\Omega$
- Good predictions of n -knockout from ^{10}Be and ^{10}C (NSCL)

Macfarlane & Pieper, PTOLEMY, ANL-76-11, Rev. 1 (1978)

Wuosmaa *et al.*, PRL **94**, 082502 (2005) + ...

Kanungo *et al.*, PLB **660**, 26 (2008)

Grinyer *et al.*, PRL **106**, 162502 (2011) + ...



ONE-NUCLEON OVERLAPS IN VMC/GFMC

For antisymmetric and translationally invariant parent $\Psi_A(\alpha)$ and daughter $\Psi_{A-1}(\gamma)$ wave functions, with $\alpha \equiv [J_A^\pi, T_A, T_{z_A}]$, $\gamma \equiv [J_{A-1}^\pi, T_{A-1}, T_{z_{A-1}}]$, and single-nucleon quantum numbers $\nu \equiv [l, s, j, t, t_z]$, the translationally invariant overlap function is:

$$R(\alpha, \gamma, \nu; r) = \sqrt{A} \left\langle [\Psi_{A-1}(\gamma) \otimes \mathcal{Y}(\nu)(\hat{r}')]_{J_A, T_A} \left| \frac{\delta(r - r')}{r^2} \right| \Psi_A(\alpha) \right\rangle$$

where $\mathcal{Y}(\nu)(\hat{r}') = [Y_l(\hat{r}') \otimes \chi_s]_j \chi_t$ and $|\Psi_{A-1}(\gamma)|^2 = 1$, $|\Psi_A(\alpha)|^2 = 1$.

The corresponding spectroscopic factor is the norm of the overlap:

$$S(\alpha, \gamma, \nu) = \int |R(\alpha, \gamma, \nu; r)|^2 r^2 dr$$

Overlap functions R satisfy a one-body Schrödinger equation with appropriate source terms. Asymptotically, at $r \rightarrow \infty$, these source terms contain core-valence Coulomb interaction at most, and hence for parent states below core-valence separation thresholds:



$$R(\alpha, \gamma, \nu; r) \xrightarrow{r \rightarrow \infty} C(\alpha, \gamma, \nu) \frac{W_{-\eta, l+1/2}(2kr)}{r},$$

where $W_{-\eta, l+1/2}(2kr)$ is a Whittaker function with $k = \sqrt{2\mu B}/\hbar$, B is the separation energy, and $C(\alpha, \gamma, \nu)$ is the asymptotic normalization coefficient or **ANC**.

GFMC evaluation of R is by extrapolation requiring two mixed estimates minus the VMC result:

$$R(\alpha, \gamma, \nu; r; \tau) \approx \langle R(\alpha, \gamma, \nu; r; \tau) \rangle_{M_A} + \langle R(\alpha, \gamma, \nu; r; \tau) \rangle_{M_{A-1}} - \langle R(\alpha, \gamma, \nu; r) \rangle_V,$$

where M_A denotes a mixed estimate where parent $\Psi_A(\alpha; \tau)$ has been propagated in GFMC and M_{A-1} is a mixed estimate where daughter $\Psi_{A-1}(\gamma; \tau)$ has been propagated.



Imaginary time evolution of overlaps in the $p_{3/2}$ channel of the overlap $\langle {}^6\text{He} + p | {}^7\text{Li} \rangle$

ALTERNATE ROUTE TO ANCS

The VMC wave functions account fairly well for short-range correlations but may have poor asymptotic behavior, particularly in p-shell.

Fitting $C = rR(r)/W(2kr)$ is generally difficult because long-range shapes can be wrong, and Monte Carlo sampling of the tails is difficult.

An alternative to explicit computation of the overlap function is an integral over the wave function interior:

$$C_{lj} = \frac{2\mu}{k\hbar^2 w} \mathcal{A} \int \frac{M_{-\eta, l+\frac{1}{2}}(2kr_{cc})}{r_{cc}} \Psi_{A-1}^\dagger \chi^\dagger Y_{lm}^\dagger(\hat{\mathbf{r}}_{cc}) (U_{\text{rel}} - V_C) \Psi_A d\mathbf{R}$$

$M_{-\eta, l+\frac{1}{2}}(2kr)$ is the “other” Whittaker function, irregular at $r \rightarrow \infty$. Here U_{rel} is


$$U_{\text{rel}} = \sum_{i < A} v_{iA} + \sum_{i < j < A} V_{ijA}$$

and at large separation of the last nucleon, $U_{\text{rel}} \rightarrow V_C$, so $(U_{\text{rel}} - V_C) \rightarrow 0$.

This makes the integrand terminate at ~ 7 fm for many p-shell nuclei.




Here is a case where fitting to VMC samples is impossible, but the integral method using the laboratory separation energy works beautifully:



ANC (fm^{-1})	VMC: AV18+UIX binding	VMC: Lab binding	Experiment
$C_{p\ 1/2}^2$	0.029(2)	0.048(3)	0.048(6)
$C_{p\ 3/2}^2$	0.237(9)	0.382(14)	0.384(38)

RESULTS FOR ONE-NUCLEON REMOVAL $3 \leq A \leq 9$



- Small error bars are VMC statistics
- Large ones are “experimental”
- Sensitivity to wave function construction seems weak but hard to quantify
- $A \leq 4$ clearly dominated by systematics, also old
- With a few exceptions, these are the first *ab initio* ANCs in $A > 4$
- $S_{17}(0) = [38.7(\text{eV b fm})] |C(2^+, {}^8\text{B})|^2 = 20.8 \text{ eV b} = \text{Solar fusion II recommended value}$
- Similar integral relation can give good estimate of excited state widths

Nollett and Wiringa, PRC **83**, 041001(R) (2011)

Nollett, arXiv:1206.0046

GFMC FOR SCATTERING STATES


GFMC treats nuclei as particle-stable system – should be good for energies of narrow resonances
Need better treatment for locations and widths of wide states and for capture reactions

METHOD

- Pick a logarithmic derivative, χ , at some large boundary radius ($R_B \approx 9$ fm)
- GFMC propagation, using method of images to preserve χ at R , finds $E(R_B, \chi)$
- Phase shift, $\delta(E)$, is function of R_B, χ, E
- Repeat for a number of χ until $\delta(E)$ is mapped out
- need E accurate to $\sim 1/3\%$

Example for ${}^5\text{He}(\frac{1}{2}^-)$

- “Bound-state” boundary condition does not give stable energy; Decaying to $n+{}^4\text{He}$ threshold
- Scattering boundary condition produces stable energy.





${}^5\text{He}$ AS $n+{}^4\text{He}$ SCATTERING

Black curves: Hale phase shifts from R -matrix analysis up to $J = \frac{9}{2}$ of data

AV18 with no V_{ijk} underbinds ${}^5\text{He}(3/2^-)$ & overbinds ${}^5\text{He}(1/2^-)$

AV18+UIX improves ${}^5\text{He}(1/2^-)$ but still too small spin-orbit splitting

AV18+IL2 reproduces locations and widths of both P -wave resonances



CONCLUSIONS

We have demonstrated that realistic nuclear Hamiltonians and accurate QMC calculations can reproduce many properties of light nuclei:

- Argonne v_{ij} + Illinois V_{ijk} gives rms binding-energy errors < 0.6 MeV for $A = 3-12$
- Successfully predict/reproduce densities, radii, moments, & transition matrix elements
- Can obtain energies and widths of low-energy nucleon-nucleus scattering states

There are many more exciting challenges in the structure and reactions of $A \leq 12$ nuclei, which we want to tackle in the next few years, such as:

- ^{12}C excited states and transitions; ν - ^{12}C scattering
- Single- & double-intruder states in $^{9,10,11}\text{Be}$, $^{10,11}\text{B}$; ^{11}Li
- More electroweak transitions in $A \leq 12$
- Charge-independence breaking in ^8Be isospin-mixing, $^{10}\text{C}(\beta^+)^{10}\text{B}$
- Parity-violating n - α scattering: $\langle ^5\text{He}(\frac{1}{2}^-) | H_{PV} | ^5\text{He}(\frac{1}{2}^+) \rangle$
- Cluster-cluster overlaps, SFs, ANCs, for $\langle (A-2)d | A \rangle$, $\langle (A-4)\alpha | A \rangle$
- Astrophysical reactions such as $^3\text{He}(\alpha, \gamma)^7\text{Be}$

For larger nuclei $A > 12$ some possibilities are:

- exascale computing for ^{16}O ($\sim 1000\times$ more expensive than ^{12}C)
- cluster GFMC (cluster VMC for ^{16}O done in 1990s)
- AFDMC (auxiliary field diffusion Monte Carlo) or hybrid GFMC-AFDMC

