

VANDERBILT
UNIVERSITY

How to measure energy-energy correlators from heavy ion collisions

Jussi Viinikainen

Vanderbilt University

Probing QCD at High Energy and Density with Jets
INT WORKSHOP INT-21R-2B

Contents

- 1 Introduction
- 2 Dealing with background
- 3 Unfolding strategies
- 4 Simple energy-loss model

Contents

- 1 Introduction
- 2 Dealing with background
- 3 Unfolding strategies
- 4 Simple energy-loss model

- A two-point energy correlator can be defined as

$$\frac{d\Sigma}{d\theta} = \int d\vec{n}_{1,2} \frac{\langle \epsilon(\vec{n}_1) \epsilon(\vec{n}_2) \rangle}{Q^2} \delta^2(\vec{n}_1 \cdot \vec{n}_2 - \cos(\theta))$$

where $\epsilon(\vec{n}_i)$ measures the asymptotic energy flux in the direction \vec{n}_i , θ is the angle between \vec{n}_1 and \vec{n}_2 , and Q is the hard scale

- Energy-energy correlators are predicted to be sensitive to several phenomena in heavy-ion collisions:
 - Color coherence: arXiv:2209.11236, arXiv:2303.03413
 - Jet wake: arXiv:2310.01500
 - Dead cone: arXiv:2307.15110
 - Anisotropic jet shapes: arXiv:2308.01294
 - ...


Experimental definition of energy-energy correlator

$$\frac{d\Sigma}{d\theta} = \int d\vec{n}_{1,2} \frac{\langle \epsilon(\vec{n}_1) \epsilon(\vec{n}_2) \rangle}{Q^2} \delta^2(\vec{n}_1 \cdot \vec{n}_2 - \cos(\theta))$$

$$EEC(\Delta r) = C_{\text{norm}} \sum_{\text{jets} \in [\rho_{T,1}, \rho_{T,2}]} \sum_{\text{pairs}} \frac{\rho_{T,i} \rho_{T,j}}{p_{T,\text{jet}}^2} \Delta r_{i,j}$$


- C_{norm} = Normalization factor
- $\rho_{T,i} \rho_{T,j}$ = Particle transverse momentum
- $p_{T,\text{jet}}$ = Jet transverse momentum
- $\Delta r_{i,j} = \sqrt{(\Delta\eta)^2 + (\Delta\varphi)^2}$ = Angular distance between particles

Energy-energy correlators in pp collisions by STAR




- Different regions of correlator explained in arXiv:2201.07800
- Free hadron region: scaling from uniformly distributed hadrons
- Transition region: break of scaling corresponding to confinement
- Quark/gluon region: perturbative interactions between quarks/gluons

Energy-energy correlators in pp collisions by ALICE



- Higher p_T jets peak at smaller opening angles between particles
 - Higher initial virtuality, more energy to radiate to reach Λ_{QCD}
 - Angular ordering \Rightarrow hadronization starts at later time!

Energy-energy correlators in pp collisions by CMS



- Shift of peak in EEC continues until very high jet p_T
- Different event generators give different predictions
- No event generator can describe data over the whole jet p_T range

Contents

- 1 Introduction
- 2 Dealing with background**
- 3 Unfolding strategies
- 4 Simple energy-loss model

Simulations used in this study

- Pythia8 (CP5)
 - Study energy-energy correlators in vacuum
- Pythia+Hydjet
 - Hard jet events generated by Pythia8 (CP5) are embedded into soft underlying event generated by Hydjet (Drum5F)
 - Select the 0-10% most central events based on UE energy density
 - Jet energy loss is not simulated
 - Jet reconstruction is done only using Pythia8 particles
 - ⇒ Particle from Pythia = signal
 - ⇒ Particle from Hydjet = background

Signal	Pythia+Pythia	Signal+Signal
Background	Pythia+Hydjet	Signal+Fake
	Hydjet+Hydjet	Fake+Fake


Energy-energy correlator definition for this analysis

$$\text{EEC}(\Delta r) = C_{\text{norm}} \sum_{\text{jets} \in [\rho_{T,1}, \rho_{T,2}]} \sum_{\text{pairs}} \frac{p_{T,i} p_{T,j}}{p_{T,\text{jet}}^2} \Delta r_{i,j}$$

$$\text{EEC}(\Delta r) = \frac{1}{N_{\text{pairs}}} \sum_{\text{jets} \in [\rho_{T,1}, \rho_{T,2}]} \sum_{\text{pairs}} p_{T,i} p_{T,j} \Delta r_{i,j}$$


- Normalize with weighted number of pairs
 - ⇒ Integral over analyzed area is one
- Do not add jet p_T to the pair weight
 - Improves resolution, no need for unfolding
- Selects pairs within $R = 0.4$ from winner-take-all jet axis

Energy-energy correlators as a function of jet p_T



- The shape of the correlators are similar to pp data
 - Regions for free hadrons, transition, and free quarks/gluons visible
 - $0.4 < \Delta r$: Acceptance drop outside of jet radius
 - Higher p_T jets peak at lower values of Δr


Energy-energy correlators as a function of track p_T




- For lower track p_T cuts, enhancement close to jet radius
- This is caused by background contributions
- Is high track p_T cut enough to suppress background?

Expected background from Pythia+Hydjet simulation

1 GeV < track p_T




3 GeV < track p_T



- Significant background contribution with 1 GeV cut
- Even with 3 GeV cut, significant background around $0.2 \lesssim \Delta r \lesssim 0.4$
- Background subtraction needed in PbPb

Background estimation: reflected η cone


- Reflect jet η coordinate, require at least twice the cone radius distance from original axis to avoid overlapping cones
 - if $|\eta_{\text{jet}}| > R \Rightarrow \eta_{\text{reflected}} = -\eta_{\text{jet}}$
 - if $-R \leq \eta_{\text{jet}} < 0 \Rightarrow \eta_{\text{reflected}} = \eta_{\text{jet}} + 2R$
 - if $0 \leq \eta_{\text{jet}} \leq R \Rightarrow \eta_{\text{reflected}} = \eta_{\text{jet}} - 2R$
- The background estimation is constructed by pairing all particles from the **signal cone** with all particles in the **reflected cone**



Normalization of reflected η cone background


- Even when one reflected cone per event is used to estimate background, this still needs to be normalized
 - Assume 10 background particles in signal cone and 10 in reflected cone
 - True background: $9 + 8 + \dots + 2 + 1 = 45$ pairings
 - Reflected cone estimate: $10 + 9 + \dots + 2 + 1 = 55$ pairings
 - There might be a random jet in the reflected cone
- ⇒ Estimate the effect on total normalization from MC by taking ratio of background pairs in signal cone and all pairs in reflected cone

Fake+fake background within reflected η cone




- Pairing reflected cone particles reproduces fake+fake contribution within signal cone

Signal+fake background estimation with reflected η cone






- Pairing signal particles from jet cone with reflected cone particles reproduces signal+fake contribution

Fake+fake background estimation with reflected η cone



- Pairing background particles from jet cone with reflected cone does not reproduce fake+fake contribution. Missing local correlations

Overall performance of reflected η cone




- Left: pairing reflected cone particles reproduces fake+fake contribution within signal cone
- Middle: pairing signal particles from jet cone with reflected cone particles reproduces signal+fake contribution
- Right: pairing background particles from jet cone with reflected cone does not reproduce fake+fake contribution. Missing local correlations.


⇒ Restrict analysis to region where fake+fake contribution is small

Determine region of validity for reflected η cone


track $p_T > 0.7$ GeV




track $p_T > 1$ GeV




track $p_T > 1.5$ GeV




track $p_T > 2$ GeV



track $p_T > 2.5$ GeV



track $p_T > 3$ GeV




- Better than 5% closure at $p_T > 2$ GeV and 1% at $p_T > 2.5$ GeV

Contents

- 1 Introduction
- 2 Dealing with background
- 3 Unfolding strategies**
- 4 Simple energy-loss model

Unfolding the measurement, how is this done in pp?


- Example from CMS-PAS-SMP-22-015
- Three-dimensional unfolding with variables:
 - Jet p_T
 - Energy weight $\frac{p_{T,i} p_{T,j}}{p_{T,\text{jet}}^2}$
 - Pair opening angle Δr




- Multidimensional unfolding in heavy ion collisions is difficult
 - Worse jet resolution compared to pp
 - Use of vacuum reference questionable
- Is there a way to simplify the unfolding procedure?

Does my observable need unfolding: jet p_T

- Jet p_T has significant resolution effects
- Gaussian smearing with $\sigma = 0.16$ applied for jet p_T to estimate resolution effects
- Resolution combined with steeply falling spectrum \Rightarrow correlator shifts to right
- Conclusion: jet p_T needs unfolding




Does my observable need unfolding: Δr




- Resolution model: certain probability Δr moves to neighboring bin

- Particle pair Δr resolution effects only important at very small Δr
- Conclusion: when very small angles are avoided, no unfolding needed




Does my observable need unfolding: Δr




- Resolution model: certain probability Δr moves to neighboring bin

- Particle pair Δr resolution effects only important at very small Δr
- Conclusion: when very small angles are avoided, no unfolding needed




Does my observable need unfolding: $p_{T,1} p_{T,2}$ weight

- Particle p_T resolution generally good
- Gaussian smearing with $\sigma = 0.024$ applied for $p_{T,1} p_{T,2}$ to estimate resolution effects
- Conclusion: no unfolding needed



Unfolding jet p_T as binning variable

- Unfolding with RooUnfold
 - Since unfolding deals with distributions, need to combine jet p_T bins and Δr axis
 - Create response matrix from matched reconstructed and generator level jets
 - Use generator level particles for both $\Rightarrow \Delta r$ axis is diagonal




- This construction allows unfolding jet p_T bins without touching the other variables for which resolution effects are small


Contents

- 1 Introduction
- 2 Dealing with background
- 3 Unfolding strategies
- 4 Simple energy-loss model**

Simple energy loss model: p_T spectrum shift in Pythia8




ATLAS: PRL 114 (2015) 072302
CMS: PRC 96 (2017) 015202



● Estimating energy loss effects in data

- Shift the jet p_T spectrum in Pythia8
- Find a shift that produces measured jet R_{AA} around $p_T = 120 \text{ GeV}$
- Compare energy-energy correlators in shifted and reference p_T bins

Estimate effect of energy loss to energy-energy correlator



- Comparing distribution with higher initial jet p_T to a lower one leads to narrowing of energy-energy correlator

Summary

- Energy-energy correlators have been measured in pp collisions
- Main issues in heavy ion measurement with respect to pp
 - Large background, cannot get rid of with p_T cut
 - Worse object resolution, more challenging unfolding
- Methods exist to mitigate these effects
 - Reflected η cone background estimate + p_T cut
 - Clever weight definition and binning choices reduce dimensions to unfold
- Bottom line: there is a way to measure energy-energy correlators in heavy ion collisions! Stay tuned for results!

This work is supported by the grant DE-FG05-92ER40712 from the US Department of Energy





Image credit: BOOST 2022 conference logo

Particle density with respect to jet axis in Pythia8

- E-Scheme axis has a dip in particle density around jet radius
- In correlation measurements, good to avoid sharp structures like this



Energy-energy correlator axis comparison in Pythia8

- Most of the pairs are the same
- For e-scheme axis, strong enhancement with respect to WTA around the jet radius

