


Data-driven Constraints on NME Calculations for Mo-100 $0\nu\beta\beta$ with CUORE and CUPID

and other topics of potential interest




Yury Kolomensky
UC Berkeley/LBNL
July 21, 2026



CUORE

Array of 988 TeO₂ crystals

- 19 towers suspended in a cylindrical structure (988 crystals)
- 5x5x5 cm³ (750g each); ¹³⁰Te: 34.1% natural isotope abundance
- **750 kg TeO₂ => 206 kg ¹³⁰Te**
- Pulse tube refrigerator and cryostat: coldest m³ in the Universe ! [arXiv:1410.1560]
- Radio-purity techniques and high resolution: low backgrounds
- Joint venture between Italy (INFN) and US (DOE, NSF) at LNGS in Italy
- Data taking since 2017, now >3 ton-years of TeO₂ exposure collected



Dell'Oro S. et al., Cryogenics 102, 9, (2019)
<https://doi.org/10.1016/j.cryogenics.2019.06.011>

Adams D. et al. (CUORE collaboration), Prog.Part.Nucl.Phys. 122 (2022) 103902,
<https://doi.org/10.1016/j.pnpnp.2021.103902>

CUPID



Next-generation bolometric ton-scale experiment at LNGS

Mission: Discover $0\nu\beta\beta$ if $m_{\beta\beta} > 10$ meV (half-life in $^{100}\text{Mo} > 10^{27}$ years)





R. Artusa et al., Eur.Phys.J. **C74**, 3096 (2014)
pre-CDR: arXiv:1907.09376

- Baseline: 1500 enriched Li_2MoO_4 crystals (~ 240 kg of ^{100}Mo)
- Demonstrated radio-purity, active background rejection
- Energy resolution ~ 5 keV
- Total background of < 0.1 counts/(ton*kev*year)
- Phased deployment options up to 1 ton of ^{100}Mo (CUPID-1T)

Based on CUORE cryogenic technologies and Scintillating Bolometer R&D by Lucifer/CUPID-0, LUMINEU, CUPID-Mo




CUPID Demonstrators and Prototypes



CUPID-Mo: 4 kg of enriched Li_2MoO_4 crystals
 Operated 2019-2020 at LSM
 1.48 kg $\times$ yr of ^{100}Mo exposure


CROSS: 9 kg of enriched Li_2MoO_4 crystals (5 kg of ^{100}Mo) + 3.3 kg of enriched TeO_2 (2.4 kg of ^{100}Te). Open structure. Currently operating at LSC



Measurements Relevant to $0\nu\beta\beta$ NMEs

- Closed-packed bolometric detectors: high statistics, low energy thresholds, high efficiency, some spatial segmentation
- Measurements (possibly) relevant for NME calculations:
 - Precision measurements of $2\nu\beta\beta$ half-life **and** spectral shape
 - $2\nu\beta\beta$ decays to excited states of the daughter nuclei
 - Possibly $2\nu\beta\beta$ in multi-site events, sensitive to the energy distribution of individual electrons

Measurement of $2\nu\beta\beta$ half-life in ^{100}Mo in CUPID-Mo




$$T_{1/2}^{2\nu} = (7.07 \pm 0.02 \text{ (stat.)} \pm 0.11 \text{ (syst.)}) \times 10^{18} \text{ yr}$$

One of the most precise measurements of $T_{1/2}$ in any isotope

- Precision measurement of half-life and spectral shape
- High statistics, low background
- Detailed background model fit with 41 sources at various locations in the detector
 - ➡ ^{238}U , ^{232}Th chains, ^{40}K , ^{90}Sr , cosmogenic and neutron-induced backgrounds
 - ➡ Constrain backgrounds by simultaneous fit to single- and multi-site, β/γ and α spectra
 - ➡ Signal efficiency constrained by the measured activity of a ^{56}Co calibration source

CUPID-Mo Background Model



Measurement of the $2\nu\beta\beta$ Spectrum in ^{100}Mo

- High statistics and low backgrounds allow constraints on the spectral shape of the $2\nu\beta\beta$ spectrum

- ☞ Sensitivity to the interplay between low- and high-energy excitations in the $2\nu\beta\beta$ transitions
- ☞ Taylor expansion of the energy spectrum in terms of spectral functions $G(E)$ and “shape parameters” ξ sensitive to NME ratios

$$\frac{d\Gamma}{dE} = g_{A,\text{eff}}^4 |M_{GT-1}|^2 \left(\frac{dG_0}{dE} + \xi_{3,1} \frac{dG_2}{dE} + \frac{1}{3} \xi_{3,1}^2 \frac{dG_{22}}{dE} + \left(\frac{1}{3} \xi_{3,1}^2 + \xi_{5,1} \right) \frac{dG_4}{dE} \right)$$

$$M_{GT-1}^{2\nu} \equiv M_{GT}^{2\nu}$$



$$\xi_{31}^{2\nu} = \frac{M_{GT-3}^{2\nu}}{M_{GT-1}^{2\nu}}, \quad \xi_{51}^{2\nu} = \frac{M_{GT-5}^{2\nu}}{M_{GT-1}^{2\nu}}$$

$$M_{GT-3}^{2\nu} = \sum_n M_n \frac{4 m_e^3}{(E_n - (E_i + E_f)/2)^3},$$

$$M_{GT-5}^{2\nu} = \sum_n M_n \frac{16 m_e^5}{(E_n - (E_i + E_f)/2)^5}.$$

F. Šimkovic, R. Dvornický, D. Stefánik, A. Faessler, Phys.Rev.C **97**, 034315 (2018)


Measurement of the $2\nu\beta\beta$ Spectrum



$\xi_{3,1}$ and $\xi_{5,1}$ highly correlated: fix the ratio to the prediction of each model

$g_{A,eff}$ constraint from the combination of half-life and $\xi_{3,1}$


$$g_{A,eff}^4 = \frac{T_{1/2}^{-1} \times \xi_{3,1}^2}{M_{GT-3}^2 G}$$



$$\xi_{3,1} = 0.45 \pm 0.03 \text{ (stat.)} \pm 0.05 \text{ (syst.)} \quad \text{CUPID-Mo, PRL } \mathbf{131}, 162501 \text{ (2023)}$$

Inconsistent with HSD model, consistent with SSD (1.4σ), ISM [L. Coraggio et al., Phys. Rev. C **105**, 034312 (2022)] prediction disfavored at 2.1σ

Measurement of $2\nu\beta\beta$ half-life in ^{130}Te in CUORE




$$T_{1/2}^{2\nu} = (9.32^{+0.05}_{-0.04} \text{ stat. } ^{+0.07}_{-0.07} \text{ syst.}) \times 10^{20} \text{ yr.}$$

The most precise measurements of $T_{1/2}$ in any isotope

- Precision measurement of half-life and spectral shape
- High statistics, low-ish background
- Detailed background model fit
 - ☞ ^{238}U , ^{232}Th chains, ^{40}K , cosmogenic backgrounds
 - ☞ Constrain backgrounds by simultaneous fit to single- and multi-site spectra

Measurement of the $2\nu\beta\beta$ Spectrum in ^{130}Te





Results consistent with nuclear models

CUORE, PRL **135**, 082501 (2025)




^{100}Mo $2\nu\beta\beta$ Decay to Excited States of ^{100}Ru

- The only $2\nu\beta\beta$ transition to 0^+ and 2^+ excited state measured so far
- Precision measurement of half-life provides sensitivity to nuclear structure and quantitative cross-check of NME models
- Segmented bolometers allow detection of high-multiplicity events with high efficiency and purity




^{100}Mo $2\nu\beta\beta$ Decay to Excited States of ^{100}Ru

Category 3

 $2\nu\beta\beta$ to 0_1^+ 

540 keV peak



 $2\nu\beta\beta$ to 2_1^+ 

591 keV peak

$$T_{1/2}^{2\nu \rightarrow 0_1^+} = (7.5 \pm 0.8 \text{ (stat.) } {}^{+0.4}_{-0.3} \text{ (syst.)}) \times 10^{20} \text{ yr}$$

$$T_{1/2}^{2\nu \rightarrow 2_1^+} > 4.4 \times 10^{21} \text{ yr (90\% c.i.)}$$

$$M_{2\nu \rightarrow 0_1^+}^{\text{eff}} = 0.143 \pm 0.008 \text{ (stat.+syst.)}$$





No shape sensitivity (yet) to distinguish NME models
But the models should be able to predict the matrix element / half-life



CUPID-Mo, Phys.Rev.C **107**, 025503 (2023)

$0\nu\beta\beta$ in High-Multiplicity Events


S. Wagaarachchi
T. Zhu



Schematic of a CUORE calorimeter



Energy sum [keV/sqrt(2)]



Energy difference [keV/sqrt(2)]





10-30% of events contained in multiple crystals

Opportunity for coincidence analysis to reduce backgrounds

Sensitivity to event topology in case of discovery

$0\nu\beta\beta$ in High-Multiplicity Events

S. Wagaarachchi
T. Zhu




Fitted $0\nu\beta\beta$ rate:

$$\Gamma^{0\nu} = 3.5_{-1.8}^{+1.9} \times 10^{-25} \text{ yr}^{-1}$$

90% C.I. on ^{130}Te $0\nu\beta\beta$ half-life:

$$\tau_{1/2}^{0\nu} > 1.1 \times 10^{24} \text{ yr.}$$

CUORE Preliminary



CUPID event topologies

Holy Grail: Quantum (Phonon) TPC

- Normal ordering scale experiment will require zero background at ton scale
 - ❑ Not possible with passive techniques
 - ❑ Requires topological reconstruction (single-site vs multi-site events), position fiducialization, active veto for external γ s
 - ❑ While keeping excellent energy resolution
 - ☞ Phonon TPC: position-sensitive bolometers
 - ☞ Prototypes: DEMETER, CERES (Berkeley), OPOSSUM (LNGS)
 - ☞ External γ veto: BINGO
- Topological information may provide sensitivity to models ($0\nu\beta\beta$ and $2\nu\beta\beta$)







Image from D. C. Hurley, J. P. Wolfe, and K. A. McCarthy. Phys. Rev. B 33, 4189 (1986)



CERES: E. Brandani et al., in preparation

Detour: $\mu^- \rightarrow e^+$ Conversion in Nucleus

• Transition

$$\mu^- + A(Z, N) \rightarrow e^+ + A(Z - 2, N + 2)$$

- Incoherent conversion ($2p \rightarrow 2n$): no coherent enhancement
- Large momentum transfer: transition to ground or excited state of the nucleus

☞ Ground state branching fraction estimated to be $\sim 41\%$

☞ Most dominant excited state is a broad Giant Dipole Resonance

• Previous best measurement: SINDRUM-II

$$R_{\mu e+} \equiv \frac{\Gamma(\mu^- + {}^{48}\text{Ti} \rightarrow e^+ + {}^{48}\text{Ca}^{\text{GS}})}{\Gamma(\mu^- + {}^{48}\text{Ti} \rightarrow \nu_\mu + {}^{48}\text{Sc}^*)} < 1.7 \times 10^{-12} \text{ (@90\% C.L.)}$$







Diagram: M. Stortini

$\mu^- \rightarrow e^+$ Conversion: Complementarity




Complementarity between LFV and LNV measurements

Complementarity between nuclear (DBD) and particle physics (Mu2e) measurements

Complementarity between nuclear calculations (nuclear matrix elements)

“No-lose theorem” [Dery, Gori, Grossman, Ligeti, JHEP **10**, 100 (2024)]

Mu2e @ FNAL



Mu2e @ Fermilab

Charge-symmetric spectrometer: will be able to search for both $\mu^- \rightarrow e^-$ and $\mu^- \rightarrow e^+$!

Also sensitivity to radiative muon capture (RMC) process, through both e^+ and γ detection

Detector ready to go, expect first beam in 2027 !

Radiative Muon Capture

Radiative muon capture



$\mu^- + (Z, N) \rightarrow \nu_\mu + \gamma + (Z-1, N+1) \rightarrow \nu_\mu + e^+e^- + (Z-1, N+1)$ is the dominant background in $\mu^- \rightarrow e^+$ searches

Strongly depends on nuclear structure: RMC transitions to the ground state, excited states, or with neutron or proton emission

No predictions so far and scarcely any data ! — could use theory help

RMC endpoint spectrum on ^{27}Al will be measured by Mu2e within a week of data taking: opportunity to test nuclear models in a light nucleus

Also data (PSI) exists for Al, Si, **Ca**, **Mo**, Sn, Pb [M. Stortini, Ph.D. thesis, Yale, 2026; A. Trumic, UC Berkeley]




Transverse Asymmetries in Parity Violation Experiments

Beam-Normal Asymmetry in elastic electron-nucleus scattering

👉 **Electron beam polarized transverse to beam direction**

$$A_T \equiv \frac{2\pi}{\sigma^\uparrow + \sigma^\downarrow} \frac{d(\sigma^\uparrow - \sigma^\downarrow)}{d\phi} \propto \vec{S}_e \cdot (\vec{k}_e \times \vec{k}'_e)$$



Interference between one- and two-photon exchange


$$A_T \propto \frac{\alpha m_e}{\sqrt{s}} \approx O(ppm)$$

Potentially sensitive to nuclear structure

Purely EM process, vector couplings, no g_A quenching

Momentum transfer of $O(100 \text{ MeV})$: relevant for NLDBD ?

PREX/CREX Puzzle



Pb-208 result not currently understood

Potential interference between couplings to different excited states ?

Opportunity to bench-mark NME models ?

Additional measurements foreseen @ MAMI, MESA

Conclusions

- Spectroscopic measurements of $2\nu\beta\beta$ and other processes can be sensitive to nuclear models
- Opportunities to benchmark, tune, and possibly qualify models for the purposes of selecting $0\nu\beta\beta$ NMEs

→ Looking forward to discussing developments here !

Thanks to CUORE, CUPID, CUPID-Mo, Mu2e, E158, MOLLER collaborators

