

Hadronic Tensor with Lattice QCD

- Neutrino-Nucleon Scattering

- Neutrino-nucleon scattering via Euclidean hadronic tensor
- Inverse problem algorithms
- Elastic scattering and form factors
- Resonances and DIS


Parton Workshop, INT, Sept. 12, 2022

Hadronic tensor and neutrino-nucleus scattering

- ◆ New long-baseline neutrino experiments are in preparation: T2K, NOvA, PINGU, ORCA, Hyper-Kamiokande, DUNE...

Omar Benhar

- ◆ Be nu
- ◆ Ch cro



elastic form factors

inclusive hadronic tensor!

Parton distribution functions

inclusive hadronic tensor!

J.A. Formaggio and G.P. Zeller, RMP84, 1307 (2012)

Whitepapers on Neutrino-Nucleus Scattering

- USQCD – A. Kronfeld et al.
(Kronfeld:2019nfb)
- Snowmass – L.A. Ruso et al.
(Ruso:2022qes)

Hadronic Tensor in Euclidean Path-Integral Formalism

- Lepton Nucleon scattering
In Minkowski space

$$\frac{d^2\sigma}{dE' d\Omega} = \frac{\alpha^2}{q^4} \left(\frac{E'}{E}\right) l^{\mu\nu} W_{\mu\nu}$$



$$W_{\mu\nu}(\vec{q}, \vec{p}, \nu) = \frac{1}{\pi} \text{Im} T_{\mu\nu} = \langle N(\vec{p}) | \int \frac{d^4x}{4\pi} e^{iq \cdot x} J_\mu(x) J_\nu(0) | N(\vec{p}) \rangle_{\text{spin avg}}$$

$$= \frac{1}{2} \sum_n \int \prod_{i=1}^n \left[\frac{d^3 p_i}{(2\pi)^3 2E_{p_i}} \right] (2\pi)^3 \delta^4(p_n - p - q) \langle N(\vec{p}) | J_\mu | n \rangle \langle n | J_\nu | N(\vec{p}) \rangle_{\text{spin avg}}$$

$$W_{\mu\nu}(p, q) = -W_1(q^2, \nu) \left(g_{\mu\nu} - \frac{q_\mu q_\nu}{q^2} \right) + W_2(q^2, \nu) \left(p_\mu - \frac{p \cdot q}{q^2} q_\mu \right) \left(p_\nu - \frac{p \cdot q}{q^2} q_\nu \right)$$

- Euclidean path-integral

KFL and S.J. Dong, PRL 72, 1790 (1994)
KFL, PRD 62, 074501 (2000)



$W_{\mu\nu}$ in Euclidean Space

$$\tilde{W}_{\mu\nu}(\vec{q}, \vec{p}, \tau = t_2 - t_1) = \frac{\frac{E_P}{M_N} \text{Tr} \langle \Gamma_e \chi_N(\vec{p}, t) \sum_{\vec{x}} \frac{1}{4\pi} e^{-i\vec{q}\cdot\vec{x}} J_\mu(\vec{x}, t_2) J_\nu(0, t_1) \chi_N^\dagger(\vec{p}, 0) \rangle}{\text{Tr} \langle \Gamma_e \chi_N(\vec{p}, t) \chi_N^\dagger(\vec{p}, 0) \rangle}$$

$$\xrightarrow{t-t_2 \gg 1/\Delta E_P, t_1 \gg 1/\Delta E_P}$$

$$= \frac{1}{4\pi} \sum_n \left(\frac{2m_N}{2E_n} \right) \delta_{\vec{p}_n - \vec{p} - \vec{q}} \langle N(\vec{p}) | J_\mu | n \rangle \langle n | J_\nu | N(\vec{p}) \rangle_{\text{spin avg}} e^{-(E_n - E_P)\tau}$$

$$= \langle N(\vec{p}) | \sum_{\vec{x}} \frac{e^{-i\vec{q}\cdot\vec{x}}}{4\pi} J_\mu(\vec{x}, \tau) J_\nu(0, 0) | N(\vec{p}) \rangle_{\text{spin avg}}$$

Inverse Laplace transform – formally correct but not practical

$$W_{\mu\nu}(\vec{q}, \vec{p}, \nu) = \frac{1}{i} \int_{c-i\infty}^{c+i\infty} d\tau e^{\nu\tau} \tilde{W}_{\mu\nu}(\vec{q}, \vec{p}, \tau)$$

Laplace transform


$$\tilde{W}_{\mu\nu}(p, \vec{q}, \tau) = \int d\nu W_{\mu\nu}(p, \vec{q}, \nu) e^{-\nu\tau} \quad \longrightarrow \quad \text{inverse problem}$$

Hadronic Tensor


$$q = q_V + q_{CS}$$

Connected sea \bar{q}_{CS}

Disconnected sea $q_{DS} = (\neq ?) \bar{q}_{DS}$



Gottfried sum rule violation --
Liu and Dong, PRL 72, 1790 (1994)



Cat's ears diagrams are suppressed by $O(1/Q^2)$. $q_i^- = q_i^{v+cs} - \bar{q}_i^{cs} + q_i^{ds} - \bar{q}_i^{ds} \equiv q_i^v + q_i^{ds} - \bar{q}_i^{ds}$

Kinematics

$$\tilde{W}_{\mu\nu}(p, \vec{q}, \tau) = \frac{E_p}{m_N} \frac{\text{Tr}[\Gamma_e C_4]}{\text{Tr}[\Gamma_e C_2]} \rightarrow \sum_{\vec{x}_2 \vec{x}_1} e^{-i\vec{q} \cdot (\vec{x}_2 - \vec{x}_1)} \langle p, s | J_\mu(\vec{x}_2, t_2) J_\nu(\vec{x}_1, t_1) | p, s \rangle$$

$$\tilde{W}_{\mu\nu}(p, \vec{q}, \tau) = \int d\nu W_{\mu\nu}(p, \vec{q}, \nu) e^{-\nu\tau}$$


$$Q^2 = -q^2 = |\vec{q}|^2 - \nu^2$$

$$x = \frac{Q^2}{2m\nu}$$

$$W^2 = m^2 + 2m\nu - Q^2$$

1. Both Q^2 and ν need to be large (difficulty?)

2. Will have a range of ν (feature?)



Inverse problems are ubiquitous

- ◆ Extracting spectral functions from lattice data: $c_2(t) = \int d\omega e^{-\omega t} \rho(\omega)$

Y. Burnier and A. Rothkopf, PRL 111, 182003 (2013)

- ◆ Global fittings of PDFs: $F_i = \sum_a C_i^a \otimes f_a$

- ◆ Lattice calculation of Quasi-PDFs: $\tilde{q}(x, P_3) = \frac{2P_3}{4\pi} \sum_{z=-z_{\max}}^{z_{\max}} e^{-ixP_3z} h_{\Gamma}(P_3, z)$

J. Karpie et. al., JHEP04, 057 (2019)

$$\tilde{q}(x, \mu^2, P^z) = \int_0^1 \frac{dy}{y} Z\left(\frac{x}{y}, \frac{\mu}{P^z}\right) q(y, \mu^2) + \mathcal{O}\left(\Lambda^2/(P^z)^2, M^2/(P^z)^2\right)$$

X. Xiong et. al., PRD90:014051 (2014)

- ◆ Lattice cross sections:

$$\sigma_n(\omega, \xi^2, P^2) = \sum_a \int_{-1}^1 \frac{dx}{x} f_a(x, \mu^2) \times K_n^a(x\omega, \xi^2, x^2 P^2, \mu^2) + \mathcal{O}(\xi^2 \Lambda_{\text{QCD}}^2)$$


Y.-Q. Ma and J.-W. Qiu, PRL120, 022003 (2018)

- ◆ Lattice calculation of Pseudo-PDFs: $\mathfrak{M}_R(\nu, \mu^2) \equiv \int_0^1 dx \cos(\nu x) q_v(x, \mu^2)$

K. Orginos et al., PRD96, 094503 (2017)


Roper Puzzle

Quenched Lattice Calculations



Dynamical Fermions

M. Sun et al.
1911.02635



Visualize 2-point Functions

◆ Backus-Gilbert (BG)

G. Backus and F. Gilbert, Geophysical Journal International 16, 169 (1968)

◆ Maximum Entropy (ME)

E Rietsch et. al., JOURNAL OF GEOPHYSICS, 42:489 (1977)

M. Asakawa et. al., Prog. Part. Nucl. Phys. 46, 459 (2001)

◆ Bayesian Reconstruction (BR)

Y. Burnier and A. Rothkopf, PRL 111, 182003 (2013)

$$P[\omega | D, \alpha, m] \propto e^{Q'(\omega)} \quad Q' = \alpha S - L - \gamma(L - N_\tau)^2$$

$$S = \sum_{\nu} \left[1 - \frac{\omega(\nu)}{m(\nu)} + \log \left(\frac{\omega(\nu)}{m(\nu)} \right) \right] \Delta\nu$$


$$P[\omega | D, m] = \frac{P[D | \omega, I]}{P[D | m]} \int d\alpha P[\alpha | D, m]$$

Hyper parameter alpha is integrated over

Maximum search is in the entire parameter space ($O(10^3)$)



High precision architecture is needed (e.g., 512-bit floating point number).

Overlap fermion on 2+1 flavor DWF,
 $48^3 \times 96$, $m_\pi = 139$ MeV, $L = 5.5$ fm



RBC/UKQCD 48l, physical pion mass, $a \sim 0.11$ fm


Visualize 2-point Functions




Overlap fermion on 2+1 flavor DWF,
 $48^3 \times 96$, $m_\pi = 139$ MeV, $L = 5.5$ fm

RBC/UKQCD 48l, physical pion mass, $a \sim 0.11$ fm
Smearred grid-source, using $t=[1, 15]$, $c=1e-6$

Mock Data Test



Mock data contain two **isolated states of mass 1.0 and 1.5 GeV**, and a **constant dense spectrum (simulating the continuous spectrum)** from 2 GeV to 15 GeV. The lattice spacing is set to be 0.02 fm and the number of time slices is 100. The signal-to-noise ratio is set to be 100.





J. Liang et. al., Phys.Rev.D 101 (2020) 11, 114503

Form Factor Case for Elastic Scattering






$$\widetilde{W}_{44}(\vec{p}, \vec{q}, \tau) = \sum_{\vec{x}_2, \vec{x}_1} e^{-i\vec{q} \cdot (\vec{x}_2 - \vec{x}_1)} \langle p, s | J_4(\vec{x}_2, t_2) J_4(\vec{x}_1, t_1) | p, s \rangle = \sum_n A_n e^{-\nu_n \tau}$$

$$A_0 = \langle p, s | J_4(\vec{q}) | n = 0 \rangle \langle n = 0 | J_4(-\vec{q}) | p, s \rangle = G_E^2(Q^2)$$

RBC/UKQCD 32IF lattice (a ~ 0.063 fm, pion mass ~ 370 MeV), clover on domain wall




Higher twist contribution




X

Beyond the Elastic Contribution



u quark with elastic contribution subtracted



Non-zero inelastic contribution

Need more data with finer lattice spacings and better understanding of the reconstruction

Compare different inverse methods



Phenomenological input

$$W^2 = M_N^2 + 2M_N\nu - Q^2$$

Large Momentum Transfer Case

$\mu = \nu = 1$ and $p_1 = q_1 = 0$ $W_{11}(\nu) = F_1(x, Q^2)$
 $24^3 \times 128$, $a_s \sim 0.12$ fm, $\xi \sim 3.5$, $m_\pi \sim 380$ MeV, $\frac{2\pi}{L} \sim 0.42$ GeV

p	q	E_p	$E_{n=0}$	$ q $	ν	Q^2	x
(0,3,3)	(0,-6,-6)	2.15	2.15	3.57	[2.96, 3.28]	[4, 2]	[0.16, 0.07]



J. Liang et al., Phys.Rev.D 101 (2020) 11, 114503

Fine lattice spacings!
 Better solving the inverse problem!

$$W^2 = (p + q)^2 = M_N^2 - Q^2 + 2E_p\nu - 2\vec{p} \cdot \vec{q}$$

$$Q^2 = 11.7 \text{ GeV}^2$$

$$W = 2.5 \text{ GeV}$$

$$|\vec{p} + \vec{q}| = 1.8 \text{ GeV}$$

Summary and Challenges

- How high an excitation energy can be reached on today's lattice?

$$W^2 = (p + q)^2 = M_N^2 - Q^2 + 2E_p\nu - 2\vec{p} \cdot \vec{q}$$

Take $|\vec{p}| = 1 \text{ GeV}$, $|\vec{q}| = 3 \text{ GeV}$, $\nu = 2.5 \text{ GeV} \rightarrow Q^2 = 2.75 \text{ GeV}^2$, $W = 3.35 \text{ GeV}$

- DIS is perhaps reachable on lattices with $a = 0.06 \text{ fm}$ or less, but may need to include multi-hadron interpolation field (?)
- Spectral decompositions in lattice 2-, 3-, and 4-point functions of baryons have seen limited success. Need better data.
- Spectral decomposition of 2- and 3-pt functions can be improved with a larger basis of interpolation fields in a variational approach. How about 4-pt functions?
- Can calculate the combination of V_4 , V_i , A_4 , A_i to isolate different resonance contributions in neutrino-nucleon scattering.