



# The effect of dark matter on compact stars and constraints we put on strongly interacting matter at high densities

Violetta Sagun

University of Coimbra, Portugal



In collaboration with T. Dietrich, M. Emma, E. Giangrandi, O. Ivanytskyi, I. Lopes, F. Pannarale, C. Providência, F. Schianchi



- Accumulation of DM in stars
- Effect of DM on NS properties
- Mass and Radius
- Tidal deformability and waveform
- NS cooling and heating
- Fermionic DM
- Bosonic DM
- Conclusions

Accumulation  
of DM in stars

Effect of DM  
on NS  
properties

Mass and Radius  
Tidal  
deformability and  
waveform  
NS cooling and  
heating

Fermionic DM

Bosonic DM

Conclusions

## 1 Accumulation of DM in stars

## 2 Effect of DM on NS properties

- Mass and Radius
- Tidal deformability and waveform
- NS cooling and heating

Accumulation  
of DM in stars

Effect of DM  
on NS  
properties

Mass and Radius  
Tidal  
deformability and  
waveform

NS cooling and  
heating

Fermionic DM

Bosonic DM

Conclusions

- 1 Accumulation of DM in stars
- 2 Effect of DM on NS properties
  - Mass and Radius
  - Tidal deformability and waveform
  - NS cooling and heating
- 3 Fermionic DM

Accumulation  
of DM in stars

Effect of DM  
on NS  
properties

Mass and Radius  
Tidal  
deformability and  
waveform

NS cooling and  
heating

Fermionic DM

Bosonic DM

Conclusions

- 1 Accumulation of DM in stars
- 2 Effect of DM on NS properties
  - Mass and Radius
  - Tidal deformability and waveform
  - NS cooling and heating
- 3 Fermionic DM
- 4 Bosonic DM

Accumulation  
of DM in stars

Effect of DM  
on NS  
properties

Mass and Radius  
Tidal  
deformability and  
waveform

NS cooling and  
heating

Fermionic DM

Bosonic DM

Conclusions

- 1 Accumulation of DM in stars
- 2 Effect of DM on NS properties
  - Mass and Radius
  - Tidal deformability and waveform
  - NS cooling and heating
- 3 Fermionic DM
- 4 Bosonic DM
- 5 Conclusions

Accumulation  
of DM in stars

Effect of DM  
on NS  
properties

Mass and Radius  
Tidal  
deformability and  
waveform

NS cooling and  
heating


Fermionic DM

Bosonic DM

Conclusions

- 1 Accumulation of DM in stars
- 2 Effect of DM on NS properties
  - Mass and Radius
  - Tidal deformability and waveform
  - NS cooling and heating
- 3 Fermionic DM
- 4 Bosonic DM
- 5 Conclusions

# Strongly Interacting Matter Phase Diagram



Accumulation of DM in stars  
 Effect of DM on NS properties  
 Mass and Radius  
 Tidal deformability and waveform  
 NS cooling and heating  
 Fermionic DM  
 Bosonic DM  
 Conclusions



# Constraints on the EoS

## HEP

### - proton flow

anisotropic expansion is caused by gradient of pressure, which gives an access to EoS

F. Donato et al., Science 296, 1333 (2002)



### - hadron multiplicities

hard core radii of hadrons control the rate of their production in thermal medium:  $R \approx 0.3 - 0.5$  fm

A. Andronic et al., Nucl. Phys. A 772, 187 (2006)

K. A. Bugay et al., Nucl. Phys. A 970, 133 (2018)



### - nucleon-nucleon scattering

hard core radius of nucleons extracted as a parameter of microscopic interaction potential:  $R = 0.5$  fm

M. Naghibi, Phys. Part. Nucl. 5, 924 (2014)

## Nucl. Phys.

### - nuclear matter ground state

- binding energy per nucleon at saturation density  $n_0$ :  
 $n_0 = 0.16 \pm 0.01 \text{ fm}^{-3}$ ,  $E(n_0)/A = -16.0 \pm 1.0 \text{ MeV}$
- incompressibility at  $n_0$ :

$$K_0 = 200 - 260 \text{ MeV}$$

- symmetry energy at  $n_0$ :  
 $S(n_0) = J = 30 \pm 4 \text{ MeV}$

- symmetry energy slope at  $n_0$ :  $L \equiv 3n_0 \left( \frac{\partial S(n_B)}{\partial n_B} \right)_{n_B=n_0} = 20 - 115 \text{ MeV}$

E. Khan, Phys. Rev. C, 80, 011307 (2009)

M. Dutra et al., Phys. Rev. C, 85, 032801 (2012)

Zhang, Z., Chen, L.-W., Phys. Lett. B, 726, 234 (2013)



## Chiral effective theory

- up to  $\sim 1.1 n_0$

Drischler, et al. PRC 102, 054315 (2020)  
Tews, et al. APJ 860, 149 (2018)

## Perturbative QCD

- from  $\sim 40 n_0$

Komoltsev & Kurkela, PRL 128 (2022)

## Astro

### - $\sim 2 M_{\text{sun}}$

PSR J0348+0432:  $M = 2.01^{+0.04}_{-0.04} M_{\odot}$

J. Antoniadis et al., Science, 340, 448 (2013)

PSR J0740+6620:  $M = 2.08^{+0.07}_{-0.07} M_{\odot}$

E. Fonseca et al., APJ, 915, L12 (2021)

PSR J1810+1744:  $M = 2.13^{+0.04}_{-0.04} M_{\odot}$

R. W. Ross et al., APJ, 900, L48 (2021)

PSR J0952-0607:  $M = 2.35^{+0.17}_{-0.17} M_{\odot}$


R. W. Ross et al., APJ, 934, 2 (2022)

### - NICER results

### - NSs cooling



## M-R relation



E. Ozel, F. Freire, A&A, 54, 401 (2009)

## Grav. Phys.


### GW170817 + kilonova

### NS+NS and NS+BH mergers



Love numbers and tidal polarizability are highly sensitive to EoS

LIGO and Virgo collaborations, PRL 119, 161101 (2017)  
LIGO and Virgo collaborations, arXiv:2001.01761 (2019)



## General Requirements

- causality
- thermodynamic consistency
- multicomponent character (n, p, e, ...)

Accumulation of DM in stars

Effect of DM on NS properties

Mass and Radius

Tidal deformability and waveform

NS cooling and heating

Fermionic DM

Bosonic DM

Conclusions



# DM candidates

Accumulation of DM in stars


Effect of DM on NS properties

Mass and Radius  
 Tidal deformability and waveform  
 NS cooling and heating

Fermionic DM

Bosonic DM

Conclusions



credits: Symmetry magazine

# DM accumulation regimes

- **Progenitor**

During the star formation stage the initial mixture of DM and BM contracting to form the progenitor star. Trapped DM undergoes scattering processes with baryons leading to its kinetic energy loss and thermalisation.

- **Main sequence (MS) star**

From this stage of star evolution accretion rate increases due to big gravitational potential of the star. In the most central Galaxy region  $M_{acc} \approx 10^{-5} M_{\odot} - 10^{-9} M_{\odot}$ .

- **Supernova explosion & formation of a proto-NS**

The newly-born NS should be surrounded by the dense cloud of DM particles with the temperature and radius that corresponds to the last stage of MS star evolution, i.e. a star with a silicone core.


**Kouvaris & Tinyakov 2010**

In addition, a significant amount of DM can be produced during the supernova explosion and mostly remain trapped inside the star.

- **Equilibrated NS**

$$M_{acc} \approx 10^{-14} \left( \frac{\rho_{\chi}}{0.3 \frac{\text{GeV}}{\text{cm}^3}} \right) \left( \frac{\sigma_{\chi n}}{10^{-45} \text{cm}^2} \right) \left( \frac{t}{\text{Gyr}} \right) M_{\odot}, \quad (1)$$

In the most central Galaxy region  $M_{acc} \approx 10^{-5} M_{\odot} - 10^{-8} M_{\odot}$ .



# DM and NS structure

Accumulation  
of DM in stars

Effect of DM  
on NS  
properties

Mass and Radius  
Tidal  
deformability and  
waveform

NS cooling and  
heating


Fermionic DM

Bosonic DM


Conclusions



**dark matter core**



**dark core inside a NS**



**dark halo around a NS**

Dark matter and baryon components do not expel each other but overlap due to absence of non-gravitational interaction

# Effect of DM on Mass and Radius

Accumulation of DM in stars

Effect of DM on NS properties

Mass and Radius

Tidal deformability and waveform

NS cooling and heating


Fermionic DM

Bosonic DM

Conclusions

- **DM core**  $\Rightarrow$  decrease of the maximum mass and observed stellar radius
- **DM halo**  $\Rightarrow$  increase of the maximum mass and the outermost radius

Ciarcelluti & Sandin 2011; Nelson+ 2019; Deliyergiyev+ 2019; Ivanytskyi+2020; Das+ 2020; Del Popolo+ 2020; Karkevandi+ 2022



DM core contributing to 5% of the total NS mass

$$\sqrt{\sigma_D}/m_D^3 = 0.05 \text{ GeV}^{-2}$$

Ellis+ 2018

# TOV equations - two fluid system

2 TOV equations:

$$\frac{dp_B}{dr} = - \frac{(\epsilon_B + p_B)(M + 4\pi r^3 p)}{r^2 (1 - 2M/r)}$$

$$\frac{dp_D}{dr} = - \frac{(\epsilon_D + p_D)(M + 4\pi r^3 p)}{r^2 (1 - 2M/r)}$$

BM and DM are coupled only through gravity, and their energy-momentum tensors are conserved separately

**total pressure**  $\rho(r) = p_B(r) + p_D(r)$

**gravitational mass**  $M(r) = M_B(r) + M_D(r)$ , where  $M_j(r) = 4\pi \int_0^r \epsilon_j(r') r'^2 dr'$  (j=B,D)

$M_T = M_B(R_B) + M_D(R_D)$  - total gravitational mass

Fraction of DM inside the star:

$$f_x = \frac{M_D(R_D)}{M_T}$$

# Tidal deformabilities of DM-admixed NS

Accumulation of DM in stars

Effect of DM on NS properties

Mass and Radius


Tidal deformability and waveform

NS cooling and heating

Fermionic DM

Bosonic DM

Conclusions



Tidal deformability parameter

$$\Lambda = \frac{2}{3} k_2 \left( \frac{R_{\text{outermost}}}{M_{\text{tot}}} \right)^5$$

$k_2$  – Love's number.

- $R_{\text{outermost}} = R_B \geq R_D$  - DM core
- $R_{\text{outermost}} = R_D > R_B$  - DM halo

Speed of sound should be calculated for two-fluid system **Das+ 2020**

Ellis+ 2018; Bezares+ 2019, Sagun+ 2022; Karkevandi+2022;  
Miao+2022; Leung+2022

# Effect of DM on GW waveform

Accumulation of DM in stars

Effect of DM on NS properties

Mass and Radius


Tidal deformability and waveform

NS cooling and heating

Fermionic DM

Bosonic DM

Conclusions



Guidice+ 2016; Ellis+ 2018; Bezares+ 2019

The DM cores may produce a supplementary peak in the characteristic GW spectrum of NS mergers, which can be clearly distinguished from the features induced by the baryon component

## Two-fluid 3D simulations of coalescing binary NS systems admixed with DM


**DM component:** Mirror DM (mirrors the BM to a parallel hidden sector, the same particle physics as the observable world and couples to the latter through gravity)

Berezhiani 2004; Ciancarella+ 2021

**BM component:** SLy EoS

Initial configurations

	$M_{A,B} (M_{\odot})$	Mirror dark matter %	$\rho_c^b [\rho_{\text{nuc}}]$	$\rho_c^{\text{DM}} [\rho_{\text{nuc}}]$	$R_{A,B} [\text{km}]$
SLy_M14_0	1.4	0%	3.866	0	11.45
SLy_M14_5	1.4	5%	4.360	2.234	11.00
SLy_M14_10	1.4	10%	4.713	2.854	10.60
SLy_M13_0	1.3	0%	3.624	0	11.46
SLy_M13_5	1.3	5%	4.058	2.087	11.04
SLy_M13_10	1.3	10%	4.366	2.679	10.63
SLy_M12_0	1.2	0%	3.398	0	11.46
SLy_M12_5	1.2	5%	3.791	1.960	11.04
SLy_M12_10	1.2	10%	4.056	2.499	10.65



Emma+ 2022

- higher DM fraction  $\Rightarrow$  a longer inspiral likely due to a lower deformability of dark matter admixed neutron stars.

Accumulation of DM in stars

Effect of DM on NS properties

Mass and Radius Tidal deformability and waveform

NS cooling and heating


Fermionic DM

Bosonic DM

Conclusions



# Gravitational waveform and frequency



- decrease of the disk mass  $\Rightarrow$  increasing DM fraction
- higher DM fraction  $\Rightarrow$  faster formation of the BH after the merger and harder to eject material from the bulk of the stars prior to the BH formation.
- lack of DM ejecta and debris disks  $\Rightarrow$  is related to its concentration in the NS core

	$M_{ej}$ sphere ( $M_{\odot}$ )	$M_{ej}$ integral ( $M_{\odot}$ )	$M_{disk}$ ( $M_{\odot}$ )	$f_{merger}$ [Hz]
Sly_M14_0	-	-	0.001	1770
Sly_M14_5	-	-	0.0008	2030
Sly_M14_10	-	-	0.0014	2058
Sly_M13_0	0.0168	$4.8 \cdot 10^{-3}$	0.062	1817
Sly_M13_5	0	$0.7 \cdot 10^{-3}$	0.001	1910
Sly_M13_10	0	$0.8 \cdot 10^{-3}$	0.0006	2221
Sly_M12_0	0	$0.3 \cdot 10^{-3}$	0.19*	1746
Sly_M12_5	0.0016	$2.6 \cdot 10^{-3}$	0.16*	1818
Sly_M12_10	0.0027	$3.3 \cdot 10^{-3}$	0.017	2198

## Dark matter

particle-antiparticle asymmetric



accumulated inside a star



- DM particles are fermions -> the Pauli blocking may prevent them from collapsing into a black hole

- DM particles are bosons -> at zero temperature could form Bose-Einstein condensate leading to gravitational collapse of the bosonic DM leading to the formation of a black hole

**Models of asymmetric DM should allow old NSs to exist**

Kouvaris 2013

particle-antiparticle symmetric



DM particles can annihilate



- possibility of its detection via X-ray,  $\gamma$ -ray or neutrino telescopes

Kouvaris 2008

- late-time heating -> higher surface temperature of old NSs

de Lavallaz & Fairbairn 2010  
Hamaguchi+ 2019

# Equation for thermal balance

The time evolution of the red-shifted temperature is determined by

$$C \frac{dT^\infty}{dt} = -L_\nu^\infty - L_\gamma^\infty + L_H^\infty$$

$C$  - total heat capacity of the NS


$L_\nu^\infty$  - red-shifted luminosity of the neutrino

$L_\gamma^\infty$  - red-shifted luminosity of the photon emissions


$L_H^\infty$  - red-shifted heating power

The photon emission luminosity is given by  $L_\gamma = 4\pi R^2 \sigma_B T_S^4$ , where  $\sigma_B$  is the Stefan-Boltzmann constant and  $R$  is the NS radius.

# NS cooling



Credits: Fridolin Weber



Light DM particles, such as axions, could contribute as an additional cooling channel in compact stars and their mergers

Creation mechanisms:

- nucleon bremsstrahlung
- Cooper pair breaking and formation processes

# Cooling of NS with DM

Accumulation of DM in stars

Effect of DM on NS properties


Mass and Radius  
Tidal deformability and waveform

NS cooling and heating

Fermionic DM

Bosonic DM

Conclusions




Sedrakian 2016; 2019

# Heating of NS with DM


## DM particles annihilation can cause heating of old NS

For a typical WIMP, its annihilation and capture rates equilibrate in old NSs.

### Surface Temperature



**Kouvaris 2008; Kouvaris & Tinyakov 2010;**  
**Hamaguchi+ 2019**



Evolution of the surface temperatures of a  $1.44 M_{\odot}$  neutron star situated at various galactic radii. In the present case,  $m_{\chi} = 10$  GeV,  $\sigma_0 = 1.5 \times 10^{-41}$  cm<sup>2</sup> and  $(r_{-2}, \alpha) = (16$  kpc, 0.19).

**Lavallaz & Fairbairn 2010**

## 4 NSs with mass above $2M_{\odot}$

- PSR J0348+0432:  $M = 2.01_{-0.04}^{+0.04} M_{\odot}$  (Antoniadis+ 2013)
- PSR J0740+6620:  $M = 2.08_{-0.07}^{+0.07} M_{\odot}$  (Fonseca+ 2021)
- PSR J1810+1744:  $M = 2.13_{-0.04}^{+0.04} M_{\odot}$  (Romani+ 2021)
- PSR J0952-0607:  $M = 2.35_{-0.17}^{+0.17} M_{\odot}$  (Romani+ 2022)

## Dark matter EoS

- **Asymmetric dark matter**  
relativistic Fermi gas of noninteracting particles with the spin 1/2  
**Nelson+ 2019**

## Baryon matter EoS

- **EoS with induced surface tension (IST EoS)**  
*consistent with:*  
nuclear matter ground state properties,  
proton flow data,  
heavy-ion collisions data,  
astrophysical observations,  
tidal deformability constraint from the NS-NS merger (GW170817)

VS+ 2019; VS+ 2014

Accumulation  
of DM in stars

Effect of DM  
on NS  
properties

Mass and Radius

Tidal  
deformability and  
waveform

NS cooling and  
heating

Fermionic DM

Bosonic DM

Conclusions

# Mass-Radius diagram of the DM admixed NSs

Accumulation of DM in stars


Effect of DM on NS properties

Mass and Radius  
 Tidal deformability and waveform  
 NS cooling and heating


Fermionic DM

Bosonic DM

Conclusions





$M_{max} > 2 M_{\odot}$  for any  $f_{\chi}$



for  $f_{\chi} = 3.3\%$   $M_{max}$  equals to  $2 M_{\odot}$   
 further increase of the DM fraction  
 leads to  $M_{max} < 2 M_{\odot}$



# Internal structure of the stars



$R_D = 9.4 \text{ km}$  for  $f_\chi = 0.3\%$   
 $R_D = 21.2 \text{ km}$  for  $f_\chi = 1.0 \%$   
 $R_D = 135.2 \text{ km}$  for  $f_\chi = 3.0 \%$

Large values of  $R_D$  relate to the existence of dilute and extended halos of DM around a baryon core of NS

Accumulation of DM in stars

Effect of DM on NS properties

Mass and Radius  
Tidal deformability and waveform

NS cooling and heating


Fermionic DM

Bosonic DM


Conclusions

# DM admixed NSs

### Mass-Radius diagram




### Tidal deformabilities



$$\Lambda = \frac{2}{3} k_2 \left( \frac{R_{\text{outermost}}}{M_{\text{tot}}} \right)^5 \rightarrow \Lambda(1.4M_{\odot}) < 800; \quad (2)$$

Abbott+ 2018

# Maximal mass of NS as a function of the DM fraction



for  $m_\chi = 0.174 \text{ GeV}$   $M_{max}$  is  $2 M_\odot$

DM particles with  $m_\chi \leq 0.174 \text{ GeV}$  are consistent with the  $2 M_\odot$  constraint for any  $f_\chi$

For heavier DM particles the NS mass can reach  $2 M_\odot$  only if  $f_\chi$  is limited from above

Accumulation of DM in stars

Effect of DM on NS properties

Mass and Radius

Tidal deformability and waveform


NS cooling and heating

Fermionic DM

Bosonic DM

Conclusions

# DM constraint in the Galaxy center



- $2M_{\odot}$  NS in the GC  $\Rightarrow m_{\chi} < 60$  GeV
- high DM fractions are not supported by GW170817

Measurements of M and R of compact stars at the Galaxy center will put more tight constraints on  $m_{\chi}$  and  $f_{\chi}$ .

# What is the nature of the GW190814 secondary component?



The compact binary merger event GW190814 had primary mass component, a black hole, with  $M = 23.2 M_{\odot}$  and the second component with  $M = 2.5 - 2.67 M_{\odot}$ . The nature of the secondary component raised a lot of questions.

Possible explanations:

- NS with exotic degrees of freedom, e.g. hyperons and/or quarks  
 [Tan+ 2020; Dexheimer+ 2021, Ivanyskiy+ 2022]
- highly spinning NS [Zhang & Li 2020]
- NS matter with extra stiffening of the EoS at high densities [Fattoyev+ 2020]
- BH from the 'mass gap' [Tews+ 2021; Essick & Landry 2020]

An alternative explanation, the secondary component of GW190814 is a DM-admixed NS

[Das+ 2021; Giovanni+ 2022]

Accumulation  
of DM in stars

Effect of DM  
on NS  
properties

Mass and Radius  
Tidal  
deformability and  
waveform

NS cooling and  
heating

Fermionic DM

Bosonic DM

Conclusions

# GW190814 secondary component as a dark matter admixed neutron star

Accumulation of DM in stars

Effect of DM on NS properties


Mass and Radius  
Tidal deformability and waveform

NS cooling and heating

Fermionic DM

Bosonic DM

Conclusions



Secondary component of GW190814 could be explained by the DM extended halo formation around a NS with the DM fraction  $f_\chi = 23\%$  for  $m_\chi = 100$  MeV.

# Asymmetric Bosonic Dark Matter

The minimal Lagrangian includes the complex scalar  $\chi$  and real vector  $\omega^\mu$  fields, which are coupled through the covariant derivative  $D^\mu = \partial^\mu - ig\omega^\mu$  with  $g$  being the corresponding coupling constant

$$\mathcal{L} = (D_\mu \chi)^* D^\mu \chi - m_\chi^2 \chi^* \chi - \frac{\Omega_{\mu\nu} \Omega^{\mu\nu}}{4} + \frac{m_\omega^2 \omega_\mu \omega^\mu}{2} \quad (3)$$


where  $\Omega^{\mu\nu} = \partial^\mu \omega^\nu - \partial^\nu \omega^\mu$  and  $m_\omega$  is the vector field mass.

Using a mean field approximation for  $\omega$ , we get

$$\begin{aligned}
 p_\chi &= \frac{m_I^2}{4} \left( m_\chi^2 - \mu_\chi \sqrt{2m_\chi^2 - \mu_\chi^2} \right), \\
 \varepsilon_\chi &= \frac{m_I^2}{4} \left( \frac{\mu_\chi^3}{\sqrt{2m_\chi^2 - \mu_\chi^2}} - m_\chi^2 \right),
 \end{aligned} \quad (4)$$

Chemical potential is limited

$$\begin{aligned}
 \mu_\chi &\in [m_\chi, \sqrt{2}m_\chi], \quad m_\chi - \text{boson mass} \\
 m_I &= \frac{m_\omega}{g} - \text{interaction scale}
 \end{aligned}$$



Giangrandi+ 2022 (In prep.)

# DM admixed NSs

Accumulation of DM in stars



Effect of DM on NS properties

Mass and Radius  
 Tidal deformability and waveform  
 NS cooling and heating

Fermionic DM

Bosonic DM

Conclusions





- **DM** can be accumulated in the **core** of a NS  $\Rightarrow$  significant decrease of the maximum mass and radius of a star.
- **DM halo**  $\Rightarrow$  increase of the maximum mass and the outermost radius.
- The secondary component of the GW190814 binary merger might be a DM admixed NS.

Changing the position of the NS in the Galaxy the accretion rate of DM varies, which in turn leads to different amount of DM



different modifications of  $M$ ,  $R$ ,  $\Lambda$ , surface temperature, etc

The effect of DM could mimic the properties of strongly interacting matter

Accumulation  
of DM in stars

Effect of DM  
on NS  
properties

Mass and Radius  
Tidal  
deformability and  
waveform

NS cooling and  
heating

Fermionic DM

Bosonic DM

Conclusions

# Smoking gun of the presence of DM in NSs

- **by measuring mass, radius, and moment of inertia of NSs with few-%-accuracy.**

To see this effect we need high precision measurement of  $M$  and  $R$  of compact stars as well as NS searches in the central part of the Galaxy with

**radio telescopes:** MeerKAT, SKA, ngVLA plan to increase radio pulsar timing and discover Galactic center pulsars.

**space telescopes:** NICER, ATHENA, eXTP, STROBE-X are expected to measure  $M$  and  $R$  of NSs with high accuracy.

**DM core**  $\Rightarrow$  mass and radius reduction of NSs toward the Galaxy center

**DM halo**  $\Rightarrow$  mass increase of NSs toward the Galaxy center  
or variation of mass and radius in different parts of the Galaxy

- **by performing binary numerical-relativity simulations and kilonova ejecta for DM-admixed compact stars for different DM candidates, their particle mass, interaction strength and fractions with the further comparison to GW and electromagnetic signals.**

Large statistics on NS-NS, NS-BH mergers by LIGO/Virgo/KAGRA would be very helpful  
The smoking gun of the presence of DM could be:

**supplementary peak in the characteristic GW spectrum of NS mergers; exotic waveforms; modification of the kilonova ejection;**

**post-merger regimes:** the next generation of GW detectors, i.e., the Cosmic Explorer and Einstein Telescope.

- **by detecting objects that go in contradiction with our understanding.**

As a potential candidate for a DM-admixed NS could be the secondary component of GW190814.

- **High/low surface temperature of NSs towards the Galaxy center**



# Thanks for your attention!

Accumulation  
of DM in stars

Effect of DM  
on NS  
properties

Mass and Radius  
Tidal  
deformability and  
waveform

NS cooling and  
heating

Fermionic DM

Bosonic DM

Conclusions

