

# Summary and where to go from here


## Critical points from various estimates



- YLE-1 extrapolated to continuum is at pprox (90, 700) MeV
- FSS does not give definite  $T_c$

## FRG/DSE







- FRG: estimate  $\rightarrow$  prediction?
- Influence of CP extends by 40 MeV in T and 200 MeV in  $\mu$


## YLE analysis in LQCD

- Most likely suffers from systematic uncertainties
- ullet No clear improvement path without going to lower T < 120 MeV and collecting more statistics
- ullet Tension with T' scaling observed in LQCD calculations

$$T'$$
 suggests  $\Omega(T,\mu)=\Omega(T')$ : applying to YLE,  $T\left[1+\kappa_2ig(rac{\mu_c}{T}ig)^2
ight]=T'_c=$  complex number







## (Critical) dynamics

#### Calculations in static equilibrium $\neq$ HIC:


- Finite size effects, rapid expansion.
- Non-equilibrium effects (critical slowing, memory).
- Consistent treatment of fluctuations and dissipation.


#### Static box simulations of linear size L:

- ullet Equilibration time is of order of  $au_{eq} \sim L^z$  , zpprox 3 in model H
- ullet Start from a random initial conditions: location of apparent critical point  $eq T_c$  of static equilibrium

## (Critical) dynamics

 There are two approaches to compute thermal fluctuations: Fluctuating hydro and hydro+.






• Significant progress has been made in both directions; however, it is insufficient to quantitatively determine the magnitude of CP effects on observables.


## Holes and bumps at ~20 GeV


. Flow  $\frac{dv_1}{dy}$ ,  $\frac{v_3^2}{n_{ch}}$  show minimum

HBT shows maximum

. Clusters  $\frac{tp}{d^2p}$  show peak








## Holes and bumps at ~ 20 GeV

- Energy range is in the "hydro regime"
  - Flow and HBT should come out of hydro


\_example: 
$$\frac{dv_1}{dy}$$
 by Du et al (2211.16408) by guessing at initial conditions / stopping


- . Can we do this also for HBT,  $\frac{v_3^2}{n_{ch}}$
- Cluster likely beyond hydro
  - -Spinodal? Probably not, requires phase co-existence at low μ. Also signal is very weak if phase co-existence is between QM and high density nuclear matter (Steinheimer et al)



But JAM-II also matches the data with a completely different explanation. Most models fail. Is this about baryon transport? Model space needs to be better constrained: data data data.

## Quark number scaling





Splitting between baryons and mesons is only at  $m_T - m_0 \ge 1 \, {\rm GeV}$  i.e. intermediate  $p_T$ 


To test coalescence, you must have data in that range for mesons and baryons

Coalescence at all  $p_T$ : Problems with entropy!

What is the effect of shadowing from spectators?

## Holes and bumps at ~ 20 GeV


Could it be due to the cross over transition?



What make the bumps/holes. Likely not the evolution though crossover

## Cumulants


Zachary Sweger (for STAR) QM 2025




UrQMD (WITHOUT mean field) gets energy dependence qualitatively right!!!

# The (possible) culprit

X. Zhang, Y. Zhang, X. Luo, N. Xu, arXiv: 2506.18832



Fluctuating impact parameter STAR centrality selection





Fixed impact parameter (b=3 fm) minimal volume fluctuations.


N.B.: Centrality Bin Width Corrections applied to both

Possible culprit: volume (N<sub>part</sub>) fluctuations

## Higher order cumulants







Error bars are unfortunately sizable. Theory has uncertainties as well!!!! At best "supporting evidence" Concentrate on 2 and 3 order!

## Test of baseline: Reduce correlation functions


reduced correlation functions


$$\hat{c}_k = \frac{\hat{C}_k}{\hat{C}_1^k} = \frac{\kappa_k}{\kappa_1^k} = \frac{FC_k}{FC_1^k}$$
 should be flat as a function of rapidity window

 Test with BES 2 data with AND WITHOUT CBWC corrections




- Effect may be small
- (BES I) anti protons do NOT agree with baseline!!!!!
- UrQMD (at 3 GeV) does NOT seem to give a flat curve





#### Acceptance Dependence






INT Workshop on "The QCD Critical Point Search: Are We There Yet"


X.Dong/LBNL


### Cumulants

- Need to understand the UrQMD energy dependence before we conclude about attractive/ repulsive
- Second order cumulants are integrals of 2 particle correlation functions which can/have been measured.
  - Plus side: Extra constraints
  - Possible issue: Sensitive to all kind of microscopic dynamics such as flow etc



## pt fluctuations






Differential pt correlations should help to resolve this issue

## Dileptons


- Dileptons can provide info about the T reached and thus help calibrate the calculations.
- Signal from CP likely not measurable  $m \leq 150\,\mathrm{MeV}, p_t \sim 5\,\mathrm{MeV}$
- 1st order PT may require absolute normalization of calculations. Is this possible?
- Chiral symmetry???



## Finite size scaling

- Seems to work
- gets critical chemical potential close to other approaches
- fails for lower energies
- works for Smash !?!

TEST: will it work for HRG???




## 4D fit $\nu$ , $\beta$ , $\gamma$ , $\mu_c$ for STAR data and SMASH cascade

STAR:  $\chi_2$ 

 $\nu = 0.76$ ,  $\beta = N/A$ ,  $\gamma = 1.33$ ,  $\mu_c = 622 \text{ MeV}$ 


STAR\_chi2\_scaling\_plot\_nu=0.760\_beta=0.120\_gamma=1.330\_muBc=0.622\_Nybins=5\_Ndata=9



SMASH:  $\chi_2$ 

$$\nu = 0.76$$
,  $\beta = N/A$ ,  $\gamma = 1.33$ ,  $\mu_c = 599$  MeV

SMASH\_chi2\_scaling\_plot\_nu=0.760\_beta=0.120\_gamma=1.330\_muBc=0.599\_Nybins=5\_Ndata=6



MF:  $\nu = 0.5$ ,  $\gamma = 1$ ,  $\beta = \frac{1}{2}$ ,  $\chi_2 \sim t^{-1}$ ,  $\chi_3 \sim t^{-2.5}$ , 3D Ising:  $\nu = 0.630$ ,  $\gamma = 1.237$ ,  $\beta = 0.326$ ,  $\chi_2 \sim t^{-1.237}$ ,  $\chi_3 \sim t^{-2.8}$ 

### **Neutron Stars**


Neutron star studies put constraints on the QCD EOS

Yes

What can they tell us about the CP of nearly-isospin symmetric matter?

We all agree, right now, nothing.

Well, maybe a bit?!




 $0.25 \atop \log_{10}(n/n_s)$ 

0.75


1.00


0.2



# Going forward

# Going forward Option 1





# Going forward Option 2: "Homework problems"

- Is hydro (+ afterburner) able to reproduce the bumps and holes in flow and HBT?
  - What need to change in baryon transport in dynamical models to get e.g.  $\frac{dv_1}{dv}$  etc
- Where does quark number scaling arise in a dynamical model which also reproduces all flow data? Relaxation time viscous correction effects??? (Teany et al)
  - Get the viscous correction from UrQMD and then plug into hydro (Thomas Schäfer)
  - Low energy: effect of shadowing???
- Check/verify that hydro with CP leads to same flow etc results as without CP
- BESII data for clusters
- Finite size scaling for HRG?
- BESII data for reduced correlation functions with and without CBWC
- Check if UrQMD gives constant reduced correlation functions
  - It seems not to. Why?
- Reduced correlations for anti-protons from BESII. Still a problem w.r.t baseline?

# Going forward Option 2: "Homework problems"

- FRG, DSE, AdS:
  - - $\epsilon$ ,  $n_B$ , s, p along cross-over (and possibly co-existence) line
  - Slope of co-existence line at CP, streng of singularity, angle between t and h
  - Comparison with Lattice at imaginary chemical potential
  - Transport coefficients??
  - Dynamics??????
- Lattice, FRG, DSE, AdS: Please provide (net) proton cumulants with baryon number observations (for example using using Kitazawa et al (1107.2755) and SAM (2003.13905))
- Stochastic hydro/Hydro plus:
  - Calculation of expanding system (critical and non-critical)
  - provide upper limit of signal for CP?

### Homework

- List what is already done and send plot to Agnieszka
- Systematics for Cumulant baselines

# The chef recommends



# Backup